arXiv:1506.03601v1 [math.RT] 11 Jun 2015

ISRAEL JOURNAL OF MATHEMATICS **00** (XXXX), 1–34
DOI: 10.1007/s00000000000000000000000

REPRESENTATIONS OF $D_q(\Bbbk[x])$

BY

Vyacheslav Futorny

Departamento de Matemática, Universidade de São Paulo, Caixa Postal 66281, São Paulo-SP, Brazil 05315-970.
e-mail: futorny@ime.usp.br

AND

Uma N. Iyer

Department of Mathematics and Computer Science, Bronx Community College, 2155 University Avenue, Bronx, New York 10453, USA.
e-mail: uma.iyer@bcc.cuny.edu *

ABSTRACT

The algebra of quantum differential operators on graded algebras was introduced by V. Lunts and A. Rosenberg. D. Jordan, T. McCune and the second author have identified this algebra of quantum differential operators on the polynomial algebra with coefficients in an algebraically closed field of characteristic zero. It contains the first Weyl algebra and the quantum Weyl algebra as its subalgebras. In this paper we classify irreducible weight modules over the algebra of quantum differential operators on the polynomial algebra. Some classes of indecomposable modules are constructed in the case of positive characteristic and q root of unity.

Contents

1. Introduction 2
2. Preliminaries 4
3. Indecomposable modules over A_q and A_1. 7
4. Irreducible D-modules 21
Acknowledgements 33

---

* Corresponding author.

Received on September, 2014

References 33

1. Introduction

Let $\Bbbk = \mathbb{Q}(q)$ be a field where $q \in \mathbb{C}^*$ is not a root of unity. Let $\Bbbk[x]$ denote the polynomial algebra over $\Bbbk$ in one variable x. Then following [5], the algebra of quantum differential operators on $\Bbbk[x]$, denoted by $D_q(\Bbbk[x])$, for a specific bicharacter β was constructed in [4]. In [3], the algebra $D_q(\Bbbk[x])$ was described in terms of generators and relations.

View x as a linear map on $\Bbbk[x]$ via multiplication; that is, $x(f) = xf$ for $f \in \Bbbk[x]$. For $a \in \{1, -1\}$ define $\partial_a : \Bbbk[x] \to \Bbbk[x]$ linearly by $\partial_a(x^n) = \left(\frac{q^{an}-1}{q^a-1}\right) x^{n-1}$ for $n \geq 1$ and $\partial_a(1) = 0$. Further, define $\partial_0 = \partial : \Bbbk[x] \to \Bbbk[x]$ linearly by $\partial(x^n) = nx^{n-1}$ for $n \geq 0$. Then, $D_q(\Bbbk[x])$ is the subalgebra of the algebra of $\Bbbk$-linear homomorphisms, $\mathrm{Hom}(\Bbbk[x], \Bbbk[x])$, generated by the maps $\{x, \partial_1, \partial_{-1}, \partial_0\}$ ([4]). The defining relations ([3]) among these maps are

$$\partial_a x - q^a x \partial_a = 1, \tag{1.1}$$

$$\partial_a x \partial_b = \partial_b x \partial_a, \tag{1.2}$$

$$\partial_{-1}\partial_1 = q\partial_1\partial_{-1}, \quad \textit{for } a, b \in \{1, -1, 0\}. \tag{1.3}$$

In [3], it is shown that $D_q(\Bbbk[x])$ is a left and right Noetherian simple domain of GK dimension 3 when q is transcendental over $\mathbb{Q}$.

Let $\sigma : \Bbbk[x] \to \Bbbk[x]$ be the automorphism given by $\sigma(x^n) = q^n x^n$ for $n \geq 0$. Then one can check that

$$\partial_1 x - x\partial_1 = \sigma = (q-1)x\partial_1 + 1, \tag{1.4}$$

$$\partial_{-1} x - x\partial_{-1} = \sigma^{-1} = (q^{-1}-1)x\partial_{-1} + 1, \tag{1.5}$$

$$\partial_{-1} = \sigma^{-1}\partial_1. \tag{1.6}$$

Note that $D_q(\Bbbk[x])$ is $\mathbb{Z}$-graded with $deg(x) = 1$, $deg(\sigma) = deg(\sigma^{-1}) = 0$ and $deg(\partial) = deg(\partial_1) = -1$. Let $\tau = \partial x$. Then, we have $\tau\sigma = \sigma\tau$ and $\Bbbk[\tau, \sigma, \sigma^{-1}]$ is the subalgebra of $D_q(\Bbbk[x])$ of degree 0. Note that $\Bbbk[\tau, \sigma]$ is indeed a polynomial algebra, and $\Bbbk[\tau, \sigma, \sigma^{-1}]$ is the localization of $\Bbbk[\tau, \sigma]$ at powers of σ. Let $\alpha : \Bbbk[\tau, \sigma, \sigma^{-1}] \to \Bbbk[\tau, \sigma, \sigma^{-1}]$ be an algebra automorphism given by $\alpha(\tau) = \tau - 1$ and $\alpha(\sigma) = \dfrac{\sigma}{q}$.

By (1.1)-(1.3) and (1.4)-(1.6) we may view $D_q(\Bbbk[x])$ as an algebra generated over $\Bbbk[\tau, \sigma, \sigma^{-1}]$ by generators $\{x, \partial, \partial_1\}$ subject to the relations

$$\partial x = \tau, \quad x\partial = \tau - 1, \tag{1.7}$$

$$\partial_1 x = \frac{q\sigma - 1}{q-1}, \quad x\partial_1 = \frac{\sigma - 1}{q-1}, \tag{1.8}$$

$$\partial_1(\tau - 1) = \partial\left(\frac{\sigma-1}{q-1}\right), \tag{1.9}$$

$$x\tau = (\tau - 1)x, \quad x\sigma = \frac{\sigma}{q}x, \tag{1.10}$$

$$\partial\tau = (\tau+1)\partial, \quad \partial\sigma = q\sigma\partial, \tag{1.11}$$

$$\partial_1\tau = (\tau+1)\partial_1, \quad \partial_1\sigma = q\sigma\partial_1. \tag{1.12}$$

This motivates the following definition/notations.

Definition/Notations. Let $\Bbbk$ be an arbitrary field, $q \in \Bbbk^*$. For variables τ, σ, let $R = \Bbbk[\tau, \sigma, \sigma^{-1}]$ be the localization of the polynomial algebra $\Bbbk[\tau, \sigma]$ with respect to the multiplicative set $\{\sigma^i\}_{\imath \in \mathbb{Z}_{\geq 0}}$. Let $\alpha : R \to R$ be the automorphism given by $\alpha(\tau) = \tau - 1$ and $\alpha(\sigma) = \dfrac{\sigma}{q}$. Let D be the algebra generated over R with variables X, Y, Y_1 satisfying relations

$$YX = \tau, \quad XY = \alpha(\tau), \tag{1.13}$$

$$Y_1X = q\sigma - 1, \quad XY_1 = \alpha(q\sigma - 1), \tag{1.14}$$

$$Y_1(\tau - 1) = Y(\sigma - 1), \tag{1.15}$$

$$Xr = \alpha(r)X, \quad Yr = \alpha^{-1}(r)Y, \quad Y_1r = \alpha^{-1}(r)Y_1, \forall r \in R. \tag{1.16}$$

Our goal is to study weight modules of D along the lines of [1] and [2] using the fact that D contains two Generalized Weyl Algebras (GWA), denoted by A_q, A_1. In this paper we deal only with the problem of classifying irreducible modules when $\Bbbk$ is algebraically closed. We will treat the indecomposable ones in a subsequent paper.

We show that all irreducible modules in these cases come in three groups:

Family I: Irreducible D weight modules which are extended from irreducible A_q-weight modules (see section 3.1).
Family II: Irreducible D weight modules which are extended from irreducible A_1-weight modules (see section 3.2).

Family III: Irreducible D-weight modules which do not arise from the families I and II. These are indecomposable as A_q and as A_1 modules (see sections 3.1 and 3.2) by our main theorem,

THEOREM (1): *When* $\Bbbk$ *is algebraically closed, every irreducible* D *weight module is indecomposable as an* A_q*-module and as an* A_1*-module.*

Theorem 2, Theorem 3, and Theorem 4 are combined to

THEOREM (B): *The irreducible* D*-modules are as described in families I, II, and III.*

In section 4.4, we build a family of indecomposable D-modules which are decomposable as A_q and A_1 modules (see Theorem 5) when the characteristic of $\Bbbk$ is equal to the order of q. In this same set-up, we present,

THEOREM (6): *Let* $V = \oplus_{\mathfrak{m}\in\omega} V_{\mathfrak{m}}$ *be a finite dimensional* D *weight module with support contained in orbit* ω*. Then* $dim(V_{\mathfrak{m}}) = dim(V_{\mathfrak{n}})$ *for any* $\mathfrak{m}, \mathfrak{n} \in \omega$*.*

In the process, we have investigated which indecomposable weight modules of A_q and A_1 can be extended to D weight modules, which is of independent interest (sections 3.1 and 3.2). In particular, we note that in many cases, it is not possible to extend an indecomposable A_q-module (or an A_1-module) to a D-module.

Assume $\overline{\Bbbk} = \Bbbk$. For this reason, every irreducible polynomial which is needed to describe weight modules is linear.

2. Preliminaries

Let A_q be the R-subalgebra of D generated by X, Y_1. Then, A_q is a GWA:

$$Y_1 X = q\sigma - 1, \quad XY_1 = \alpha(q\sigma - 1), \tag{2.1}$$

$$Xr = \alpha(r)X, \quad Y_1 r = \alpha^{-1}(r)Y_1, \forall r \in R. \tag{2.2}$$

The algebra D is then generated over A_q by Y with relations

$$YX = \tau, \quad XY = \alpha(\tau), \tag{2.3}$$

$$Y_1(\tau - 1) = Y(\sigma - 1), \tag{2.4}$$

$$Yr = \alpha^{-1}(r)Y, \forall r \in R. \tag{2.5}$$

Analogously, let A_1 be the R-subalgebra of D generated by X, Y. Then A_1 is also a GWA with

$$YX = \tau, \quad XY = \alpha(\tau - 1), \tag{2.6}$$

$$Xr = \alpha(r)X, \quad Yr = \alpha^{-1}(r)Y, \forall r \in R. \tag{2.7}$$

D is generated over A_1 by Y_1 with relations

$$Y_1X = (q\sigma - 1), \quad XY_1 = \alpha(q\sigma - 1), \tag{2.8}$$

$$Y_1(\tau - 1) = Y(\sigma - 1), \tag{2.9}$$

$$Y_1r = \alpha^{-1}(r)Y_1, \forall r \in R. \tag{2.10}$$

In what follows, A could mean A_q or A_1, and $t = q\sigma - 1$ (if we study weight modules over A_q) or $t = \tau$ (if we study weight-modules over A_1).

Let $Max(R)$ denote the set of maximal ideals of R. For $\mathfrak{m} \in Max(R)$, set $M_\mathfrak{m} = \{v \in M \mid \mathfrak{m}v = 0\}$. We say that M is a weight module if $M = \sum_{\mathfrak{m} \in Max(R)} M_\mathfrak{m}$. Let support of M be $Supp(M) = \{\mathfrak{m} \in Max(R) \mid M_\mathfrak{m} \neq 0\}$. The cyclic group $< \alpha >$ generated by α acts on the set $Max(R)$, and let Ω denote the corresponding orbit set. When $\omega \in \Omega$ is infinite we call ω a *linear orbit.* When ω is finite it is called a *circular orbit.* Set an order on ω by $\mathfrak{m} < \alpha(\mathfrak{m})$.

For $\mathfrak{m} \in Max(R)$, let $K_\mathfrak{m} = R/\mathfrak{m}$ and $t_\mathfrak{m} = t + \mathfrak{m} \in K_\mathfrak{m}$. Call $\mathfrak{m}$ a *break* if $t_\mathfrak{m} = 0$. Let B be the set of all breaks, and $B_\omega = B \cap \omega$ for $\omega \in \Omega$. A *maximal break* in a linear orbit is $\mathfrak{m} \in B$ which is maximal with respect to the order $<$ defined in the preceding paragraph. For a circular orbit, chose any break to be the maximal break.

Note that [2] gives lists of indecomposable and simple weight GWA modules.

2.1. Description of indecomposable A-modules. For each orbit ω of r elements, fix a maximal ideal $\mathfrak{m}(\omega) \in \omega$ which is chosen to be a maximal break if ω has a break. In what follows, $T = Y_1$ if we are studying A_q-modules, and $T = Y$ if we are studying A_1-modules, and A denotes A_q or A_1.

Case $|\omega| = \infty, B_\omega = \emptyset$: Here, $V(\omega) = \oplus_{\mathfrak{m} \in \omega} K_\mathfrak{m}$ has an irreducible A-module structure with $X(v) = \alpha(t_\mathfrak{m} v)$ and $T(v) = \alpha^{-1}(v)$ for $v \in \Bbbk_\mathfrak{m}$. The module $V(\omega)$ is irreducible.

Case $|\omega| = \infty, B_\omega \neq \emptyset$: If $\mathfrak{m}$ is the maximal element in B_ω then set $B'_\omega = B_\omega \cup \{\alpha(\mathfrak{m})\}$. If B_ω does not have a maximal element, then $B'_\omega = B_\omega$. Let $J \subset B'_\omega$ be an interval, and $J' \subset J$ be any subset not containing the maximal element of J. Let $\mathfrak{n}_0$ be the maximal break in ω preceding all elements of J or $-\infty$ if it does not exist; let $\mathfrak{n}_1$ be the maximal element of J if it exists and is a break or $+\infty$ otherwise. For each $\mathfrak{m} \in \omega$, put $V_\mathfrak{m} = K_\mathfrak{m}$ if $\mathfrak{n}_0 < \mathfrak{m} \leq \mathfrak{n}_1$ and $V_\mathfrak{m} = 0$ otherwise. Set $V(\omega, J, J') = \oplus_{\mathfrak{m}\in\omega} V_\mathfrak{m}$, and for $v \in V_\mathfrak{m}$ set

$$X(v) = \begin{cases} \alpha(t_\mathfrak{m} v) & \textit{if } \mathfrak{m} \notin B, \\ \alpha(v) & \textit{if } \mathfrak{m} \in J', \\ 0 & \textit{otherwise}; \end{cases} \quad T(v) = \begin{cases} 0 & \textit{if } \alpha^{-1}(\mathfrak{m}) \in J' \cup \{\mathfrak{n}_0\}, \\ \alpha^{-1}(v) & \textit{otherwise.} \end{cases}$$

Case $|\omega| < \infty, B_\omega = \emptyset$: Fix any $f \in \Bbbk^*$. Let $V(\omega, f) = \oplus_{\mathfrak{m}\in\omega} \Bbbk_\mathfrak{m}$ and give it A-module structure by setting

$$X(v) = \begin{cases} \alpha(t_\mathfrak{m} v) & \textit{if } \mathfrak{m} \neq \mathfrak{m}(\omega), \\ f\alpha(t_\mathfrak{m} v) & \textit{if } \mathfrak{m} = \mathfrak{m}(\omega); \end{cases} \quad T(v) = \begin{cases} \alpha^{-1}(v) & \textit{if } \alpha^{-1}(\mathfrak{m}) \neq \mathfrak{m}(\omega), \\ \frac{1}{f}\alpha^{-1}(v) & \textit{if } \alpha^{-1}(\mathfrak{m}) = \mathfrak{m}(\omega). \end{cases}$$

The module $V(\omega, f)$ is irreducible.

Case $|\omega| < \infty, B_\omega \neq \emptyset$: Here, we receive two families of indecomposable families. Suppose $|B_\omega| = m > 0$. Then there is a one-to-one correspondence $\mathbb{Z}_m \to B_m$ with $\mathfrak{m}_i = \alpha^i(\mathfrak{m}_0)$ for some fixed break $\mathfrak{m}_0$. For any $\mathfrak{m} \in \omega$, let $j(\mathfrak{m}) \in \mathbb{Z}_m$ be the unique j such that $\mathfrak{m}_{j-1} < \mathfrak{m} \leq \mathfrak{m}_j$. Let x, y be two noncommuting variables.

Family 1. Fix $j \in \mathbb{Z}_m$ and $w = z_1 z_2 \cdots z_n$ be a word of length $n \geq 1$ where each $z_i \in \{x, y\}$. Let $e_0, e_1, \ldots, e_n$ be $n+1$ symbols. For each $\mathfrak{m} \in \omega$, let $V_\mathfrak{m}$ be a vector space over $\Bbbk_\mathfrak{m}$ with basis $\{(\mathfrak{m}, e_k) \mid k + j = j(\mathfrak{m}) \in \mathbb{Z}_m\}$. Put $V(\omega, j, w) = \oplus_{\mathfrak{m}\in\omega} V_\mathfrak{m}$ and it has an A-module structure given by

$$X(\mathfrak{m}, e_k) = \begin{cases} t_\mathfrak{m}(\alpha(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \notin B, \\ (\alpha(\mathfrak{m}), e_{k+1}) & \textit{if } \mathfrak{m} \in B_\omega \textit{ and } z_{k+1} = x, \\ 0 & \textit{otherwise}; \end{cases}$$

$$T(\mathfrak{m}, e_k) = \begin{cases} (\alpha^{-1}(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \textit{ is not a break}, \\ (\alpha^{-1}(\mathfrak{m}), e_{k-1}) & \textit{if } \mathfrak{m} \in B_\omega \textit{ and } z_k = y, \\ 0 & \textit{otherwise.} \end{cases}$$

Let $\mathfrak{m}_1$ be that break for which $n + j = j(\mathfrak{m}_1) \in \mathbb{Z}_m$. Then $X(\mathfrak{m}_1, e_n) = 0$. Similarly, let $\mathfrak{m}_0$ be that break for which $j = j(\mathfrak{m}_0) \in \mathbb{Z}_m$. Then $T(\mathfrak{m}_0, e_0) = 0$. The module $V(\omega, j, w)$ is irreducible if and only if w is the empty word.

Family 2. Let $w = z_1 z_2 \cdots z_n$ be a word whose length n is a multiple of m for $z_i \in \{x, y\}$. Let $f \in \Bbbk^*$. Consider n elements e_k where $k = 1, 2, \ldots, n$. For $\mathfrak{m} \in \omega$, let $V_\mathfrak{m}$ be a $\Bbbk_\mathfrak{m}$ vector space with basis $\{e_k \mid k \equiv j(\mathfrak{m})(mod\, m)\}$. The vector space $V(\omega, w, f) = \oplus_{\mathfrak{m}\in\omega} V_\mathfrak{m}$ has an A-module structure given by

$$X(\mathfrak{m}, e_k) = \begin{cases} t_\mathfrak{m}(\alpha(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \textit{ is not a break;} \\ (\alpha(\mathfrak{m}), e_{k+1}) & \textit{if } \mathfrak{m} \in B_\omega, k \neq n, z_{k+1} = x; \\ f(\alpha(\mathfrak{m}), e_1) & \textit{if } \mathfrak{m} \in B_\omega, k = n, z_1 = x; \\ 0 & \textit{otherwise;} \end{cases}$$

$$T(\mathfrak{m}, e_k) = \begin{cases} (\alpha^{-1}(\mathfrak{m}), e_k) & \textit{if } \alpha^{-1}(\mathfrak{m}) \textit{ is not a break;} \\ (\alpha^{-1}(\mathfrak{m}), e_{k-1}) & \textit{if } \alpha^{-1}((\mathfrak{m}) \in B_\omega, k \neq 1, z_k = y; \\ f(\alpha^{-1}(\mathfrak{m}), e_n) & \textit{if } \alpha^{-1}(\mathfrak{m}) \in B_\omega, k = 1, z_1 = y; \\ 0 & \textit{otherwise;} \end{cases}$$

The module $V(\omega, w, f)$ is irreducible if and only if $w = x^m$ or $w = y^m$.

3. Indecomposable modules over A_q and A_1.

3.1. THE LIST OF A_q-INDECOMPOSABLES. Here we go through the list of GWA-indecomposables as described in [2] and understand them as A_q-weight modules.

3.1.1. *The module* $V_q(\omega, b)$ *for* $b \in \Bbbk^* \setminus \{q^i\}_{i\in\mathbb{Z}}$. Suppose that $|\omega| = \infty$ and $B_\omega = \emptyset$. In this case, we see $V_q(\omega, b) = \oplus_{i\in\mathbb{Z}} \Bbbk v_i$ where $\{v_i\}_{i\in\mathbb{Z}}$ is a basis, and the action is defined by:

$$\begin{aligned} X(v_i) &= (q^{i+1}b - 1)v_{i+1}, & Y_1(v_i) &= v_{i-1}, \\ f(\tau, \sigma)(v_i) &= f(a + i, q^i b)v_i, & \textit{for } f &\in \Bbbk[\tau, \sigma, \sigma^{-1}]. \end{aligned}$$

Note in particular that $(\sigma - 1)(v_i) = (q^i b - 1)v_i \neq 0$ since $b \notin \{q^j\}_{j\in\mathbb{Z}}$. Thus, in this case we can extend the action of A_q on $V_q(\omega, b)$ to an action of D by setting

$$Y(v_i) = \frac{(a + i - 1)}{(q^i b - 1)} v_{i-1} \quad \forall i \in \mathbb{Z}.$$

Thus, for any $a \in \Bbbk$, we obtain an irreducible D-weight module, denoted by $V_q(\omega, b, a)$. Using appropriate scalar multiples of v_i, we visualize $V_q(\omega, b, a)$ without loss of generality as follows:



Remark 1: *When characteristic of* $\Bbbk$ *is* 0, *and* $a \in \Bbbk \setminus \mathbb{Z}$ *(respectively,* $a \in \Bbbk \setminus \mathbb{Z}_p$ *when characteristic of* $\Bbbk$ *is* $p > 0$*) then the resulting module* $V_q(\omega, b, a)$ *is an irreducible* A_1*-module. If* $a \in \mathbb{Z}$ *(respectively,* $a \in \mathbb{Z}_p$*), then the resulting module* $V_q(\omega, b, a)$ *is an indecomposable* A_1*-module.*

3.1.2. *The module* $V_q(\omega, J, J')$ *when* q *is not a root of unity.* Suppose $|\omega| = \infty$ and $B_\omega \neq \emptyset$. That is, $\omega = \{\alpha^i(\tau - a, \sigma - b)\}_{i \in \mathbb{Z}}$. Since $B = \{(\tau - c, q\sigma - 1) \mid c \in \Bbbk\}$, we have $B_\omega \neq \emptyset$ if and only if $b = q^k$ for some $k \in \mathbb{Z}$. That is, without loss of generality,

$$\omega = \{\mathfrak{m}_i = (\tau - a - i, \sigma - q^{i-1})\}_{i \in \mathbb{Z}}, \text{ and } B_\omega = \left\{\mathfrak{m}_0 = \left(\tau - a, \sigma - \frac{1}{q}\right)\right\}.$$

Set $B'_\omega = \{\mathfrak{m}_0, \mathfrak{m}_1\}$. Thus, we have options which we investigate one at a time: (1) $J = B'_\omega, J' = \{\mathfrak{m}_0\}$, $\mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \infty$. In this case, we have $V = \oplus_{i=-\infty}^{\infty} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \in \mathbb{Z}$. That is, $v_0 \in K_{\mathfrak{m}_0}$. Here,

$$X(v) = \begin{cases} v_1 & \text{if } v = v_0 \\ (q^i - 1)v_{i+1} & \text{if } v = v_i \end{cases} \qquad Y_1(v) = \begin{cases} 0 & \text{if } v = v_1 \\ v_{i-1} & \text{if } v = v_i \end{cases} \qquad \forall i \in \mathbb{Z}.$$

Further, $\sigma(v_i) = q^{i-1}v_i, \quad \tau(v_i) = (a+i)v_i \quad \forall i \in \mathbb{Z}$.

Using the fact that $YX = \tau$, we now define action of Y on $V_q(\omega, J, J')$ as ,

$$Y(v_{i+1}) = \frac{(a+i)}{(q^i - 1)} v_i \quad \forall i \neq 0, \quad Y(v_1) = av_0.$$

For every $a \in \Bbbk$, we thus have a D weight module denoted by $V_q(\omega, J, J', a)$ which is irreducible if $a \neq 0$, and indecomposable if $a = 0$. Note that if $a \in \mathbb{Z} \setminus \{0\}$ then $V_q(\omega, J, J', a)$ is not simple as either A_q module or A_1-module. But it is simple as a D-module. By considering suitable scalar multiples of v_i, without loss of generality, we visualize $V_q(\omega, J, J', a)$ as follows:



(2) $J = B'_\omega, J' = \emptyset$, $\mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \infty$. In this case again, we have $V = \oplus_{i=-\infty}^{\infty}\Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \in \mathbb{Z}$. Here,

$$X(v_i) = (q^i - 1)v_{i+1}, \quad Y_1(v_i) = v_{i-1}, \quad \sigma(v_i) = q^{i-1}v_i, \quad \tau(v_i) = (a+i)v_i, \forall i \in \mathbb{Z}.$$

Using the fact that $YX = \tau$, we may define Y on $V_q(\omega, J, J')$, but requiring that $XY(v_1) = (\tau - 1)(v_1)$ gives us $a = 0$. Thus, we have the following:

$$\tau(v_i) = iv_i; \quad Y(v_i) = \begin{cases} cv_0 & \textit{if } i = 1, \textit{ for some } c \in \Bbbk, \\ \left(\dfrac{i-1}{q^{i-1}-1}\right) v_{i-1} & \textit{if } i \neq 1. \end{cases}$$

For any fixed $c \in \Bbbk$, we have a D-module which is indecomposable, but not irreducible. Notice that if $c = 0$, then V is a decomposable A_1-module, but an indecomposable A_q-module which is reducible as an A_1^q-module.

A generalization of the above set-up: For any $c, d \in \Bbbk$, set $V_q(\omega, J, J', c, d) = \oplus_{i=-\infty}^{\infty}\Bbbk v_i$ with Here,

$$X(v_i) = (q^i - 1)v_{i+1}; \quad \sigma(v_i) = q^{i-1}v_i, \quad \tau(v_i) = iv_i \quad \forall i \in \mathbb{Z}.$$

$$Y_1(v_i) = \begin{cases} cv_0 & \textit{if } i = 1, \textit{ for some } c \in \Bbbk, \\ v_{i-1} & \textit{if } i \neq 1. \end{cases}$$

$$Y(v_i) = \begin{cases} dv_0 & \textit{if } i = 1, \textit{ for some } d \in \Bbbk, \\ \left(\dfrac{i-1}{q^{i-1}-1}\right) v_{i-1} & \textit{if } i \neq 1. \end{cases}$$

We have a D-module which is indecomposable (but not irreducible) for $(c, d) \neq (0, 0)$. If $c = 0$ and $d \neq 0$, then we have a module which is decomposable as an A_1^q-module, but indecomposable (but not simple) as an an A_1-module and hence indecomposable as a D-module. Conversely, if $c \neq 0$ and $d = 0$, then we have a module which is decomposable as an A_1-module but indecomposable (yet not simple) as an A_1^q-modue and hence indecomposable as a D-module. With a suitable change of basis, the picture is as follows:



Remark 2: *If $a \neq 0$, then for $J = B'_\omega, J' = \emptyset$, the module $V_q(\omega, J, J')$ cannot be extended to a D-module.*

(3) $J = \{\mathfrak{m}_0\}, J' = \emptyset, \mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \mathfrak{m}_0$

In this case again, we have $V_q(\omega, J, J') = \oplus_{i \leq 0} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \leq 0$. Here,

$$X(v_i) = \begin{cases} (q^i - 1)v_{i+1} & \text{for } i \leq -1, \\ 0 & \text{for } i = 0 \end{cases}; \qquad Y_1(v_i) = v_{i-1};$$

$$\sigma(v_i) = q^{i-1} v_i; \qquad \tau(v_i) = (a + i)v_i \quad \forall i \leq 0.$$

Since $\sigma - 1 \neq 0$, we can define action of Y on on $V_q(\omega, J, J')$ by letting $Y = Y_1\left(\dfrac{\tau - 1}{\sigma - 1}\right)$. This gives an irreducible D weight module denoted by $V_q(\omega, J, J', a)$. After a suitable change of basis, the picture here is



Remark 3: *When characteristic of $\Bbbk$ is 0, and $a \in \Bbbk \setminus \mathbb{Z}$ (respectively, $a \in \Bbbk \setminus \mathbb{Z}_p$ when characteristic of $\Bbbk$ is $p > 0$) then the resulting module $V_q(\omega, b, a)$ is an irreducible A_1-module. If $a \in \mathbb{Z}$ (respectively, $a \in \mathbb{Z}_p$), then the resulting module $V_q(\omega, b, a)$ is an indecomposable A_1-module.*

(4) $J = \{\mathfrak{m}_1\}, J' = \emptyset$, $\mathfrak{n}_0 = \mathfrak{m}_0, \mathfrak{n}_1 = \infty$ In this case we have $V_q(\omega, J, J') = \oplus_{i\geq 1}\Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \geq 1$. Here,

$$X(v_i) = (q^i - 1)v_{i+1}; \qquad Y_1(v_i) = \begin{cases} v_{i-1} & \textit{for } i \geq 2, \\ 0 & \textit{for } i = 1 \end{cases};$$

$$\sigma(v_i) = q^i v_i; \qquad \tau(v_i) = (a+i)v_i \quad \forall i \geq 1.$$

Since $\sigma - 1 \neq 0$ on $v_i, i \geq 2$, we can define Y on on $V_q(\omega, J, J')$ by letting $Y = Y_1\left(\dfrac{\tau - 1}{\sigma - 1}\right)$ on $v_i, i \geq 2$ and $Y(v_1) = 0$. This gives an irreducible D module we denote by $V_q(\omega, J, J', a)$. Here the picture is:



Remark 4: *When characteristic of* $\Bbbk$ *is* 0*, and* $a \in \Bbbk\backslash\mathbb{Z}$ *(respectively,* $a \in \Bbbk\backslash\mathbb{Z}_p$ *when characteristic of* $\Bbbk$ *is* $p > 0$*) then the resulting module* $V_q(\omega, b, a)$ *is an irreducible* A_1*-module. If* $a \in \mathbb{Z}$ *(respectively,* $a \in \mathbb{Z}_p$*), then the resulting module* $V_q(\omega, b, a)$ *is an indecomposable* A_1*-module.*

(5) $J = \emptyset, J' = \emptyset$, $\mathfrak{n}_0 = \mathfrak{m}_0, \mathfrak{n}_1 = \infty$. The resulting module is the same as the one above in case (4).
(6) $J = \emptyset, J' = \emptyset$, $\mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \infty$. The resulting module is the same as the one above in case (2).

3.1.3. *The module* $V_q(\omega, J, J')$ *when* q *is a root of unit of* 1*.* Note that since $|\omega| = \infty$ and q is a root of 1, this case arises when characteristic of $\Bbbk$ is 0. Since D contains $A_1(\Bbbk)$, we can extend only those A_q-modules which are infinite dimensional. In other words, we consider only those $J \subset B'_\omega$ for which $\mathfrak{n}_0 = -\infty$ or $\mathfrak{n}_1 = \infty$. We also have $b = q^k$ for some $k \in \mathbb{Z}$, $|B_\omega| = \infty$ and $B'_\omega = B_\omega$. Set $J' \subset J$ not containing the maximal element of J and we consider the indecomposable A_q-module $V_q(\omega, J, J')$. We point out that $V_q(\omega, J, J')$ is not irreducible A_q-module for any choice of J'.

Not every $V_q(\omega, J, J')$ can be extended to a D module. Using the condition $YX = \tau$ we see the following cases for any $v \neq 0, v \in V_{\mathfrak{m}}$:
Case 1: $X(v) = c\alpha(v) \neq 0$. Then, set $Y(\alpha(v)) = \frac{\tau(v)}{c}$.
Case 2: For $\mathfrak{m} \in J \setminus J'$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) = 0$. In this case set $Y(\alpha(v)) = dv$ for any $d \in \mathbb{k}$.
Case 3: For $\mathfrak{m} \in J \setminus J'$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) \neq 0$. In this case this A_q-module cannot be extended to a D-module since we require $YX(v) = \tau(v)$. In particular, since the characteristic of $\mathbb{k}$ is 0, we have that $\tau = 0$ can happen for at most one weight $\mathfrak{m}$. Thus, if $\mathfrak{m}_1, \mathfrak{m}_2 \in J \setminus J'$ are such that $X(v_1) = X(v_2) = 0$ for $v_1, v_2 \neq 0, v_1 \in \mathfrak{m}_1, v_2 \in \mathfrak{m}_1$, then this A_q-module cannot be extended to a D-module.

REMARK 5: *In case there is a D-module structure, then that structure is indecomposable D-module since it was indecomposable as an A_q-module to begin with. This module is irreducible as a D module if and only if $X \neq 0$ for any $\mathfrak{m} \neq \mathfrak{n}_1$ and Y, Y_1 are not simultaneously zero unless $\alpha^{-1}(\mathfrak{m}) = \mathfrak{n}_0$. Further, if $X \neq 0$ for any $\mathfrak{m} \neq \mathfrak{n}_1$ and $Y(v) = Y_1(w) = 0$ for nonzero $v, w, v \neq w, v, w \notin V_{\alpha(\mathfrak{n}_0)}$ then we have an irreducible D-module which is not irreducible as an A_1 or an A_q-module.*

3.1.4. *The module $V_q(\omega, f, b)$ for $b \notin \{q^i\}_{i\in\mathbb{Z}}$.* This family of A_q-decomposables arise when $|\omega| < \infty$ and $B_\omega = \emptyset$. In particular, the characteristic of $\mathbb{k} = p > 0$, and q is a root of 1 Since $B_\omega = \emptyset$, we have $\omega = \alpha^i(\mathfrak{m})$ where $\mathfrak{m} = (\tau - a, \sigma - b)$ where $a, b \in \mathbb{k}, b \neq 0, b \notin \{q^i\}_{i\in\mathbb{Z}}$. Suppose $|\omega| = r$ (therefore, $q^r = 1$ and p/r). For any $f \in \mathbb{k}^*$, let $V_q(\omega, f, b) = \oplus_{1\leq i\leq r}\mathbb{k}v_i$ be an r-dimensional vector space and give it A_q-module structure as described in section 2.1, with $A = A_q$, $T = Y_1$, and $V_q(\omega, f, b) = V(\omega, f)$ setting

$$\sigma(v_i) = q^{i-1}bv_i; \quad X(v_i) = \begin{cases} (q^i b - 1)v_{i+1} & \text{if } i < r, \\ f(b-1)v_1 & \text{if } i = r; \end{cases}$$

$$\tau(v_i) = (a + i - 1)v_i; \quad Y_1(v_i) = \begin{cases} v_{i-1} & \text{if } i > 1, \\ \frac{1}{f}v_r & \text{if } i = 1. \end{cases}$$

Since action of X is an isomorphism, we extend the action of A_q to an action of D by using $YX = \tau$ and setting

$$Y(v_{i+1}) = \begin{cases} \dfrac{a+i-1}{q^i b - 1} v_i & \textit{if } i > 1; \\ \dfrac{a+r-1}{f(b-1)} v_r & \textit{if } i = 1. \end{cases}$$

We denote the resulting irreducible D-module by $V_q(\omega, f, b, a)$. Note that the resulting A_1-module is also irreducible since X is invertible. With a change of basis we see the picture as given:



3.1.5. *The module* $V_q(\omega, j, w)$. This family of A_q-decomposables arise when $|\omega| < \infty$ and $B_\omega \neq \emptyset$ and this is Family 1 as described in section 2.1. Let $|B_\omega| = m > 0$ (that is, $b \in \{q^i\}_{i \in \mathbb{Z}}$). The one-to-one correspondence $\mathbb{Z}_m \to B_m$ and $j(\mathfrak{m}) \in \mathbb{Z}_m$ are as described in section 2.1. Let x, y be two noncommuting variables.

Fix $j \in \mathbb{Z}_m$ and $w = z_1 z_2 \cdots z_n$ be a word of length $n \geq 1$ where each $z_i \in \{x, y\}$. Let $e_0, e_1, \ldots, e_n$ be $n+1$ symbols. For each $\mathfrak{m} \in \omega$, let $V_\mathfrak{m}$ be a vector space over $\Bbbk_\mathfrak{m}$ with basis $\{(\mathfrak{m}, e_k) \mid k + j = j(\mathfrak{m}) \in \mathbb{Z}_m\}$. Put $V_q(\omega, j, w) = \oplus_{\mathfrak{m} \in \omega} V_\mathfrak{m}$ and it has an A_q-module structure given by

$$X(\mathfrak{m}, e_k) = \begin{cases} t_\mathfrak{m}(\alpha(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \notin B, \\ (\alpha(\mathfrak{m}), e_{k+1}) & \textit{if } \mathfrak{m} \in B_\omega \textit{ and } z_{k+1} = x, \\ 0 & \textit{otherwise}; \end{cases}$$

$$Y_1(\mathfrak{m}, e_k) = \begin{cases} (\alpha^{-1}(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \notin B, \\ (\alpha^{-1}(\mathfrak{m}), e_{k-1}) & \textit{if } \mathfrak{m} \in B_\omega \textit{ and } z_k = y, \\ 0 & \textit{otherwise.} \end{cases}$$

The module $V_q(\omega, j, w)$ is irreducible as an A_q-module when w is the empty word. Not every $V_q(\omega, j, w)$ can be extended to a D module. Using the condition $YX = \tau$ we see the following cases for any $v \neq 0, v \in V_{\mathfrak{m}}$:
Case 1: $X(v) = c\alpha(v) \neq 0$. Then, set $Y(\alpha(v)) = \frac{\tau(v)}{c}$.
Case 2: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) = 0$. In this case set $Y(\alpha(v)) = dv$ for any $d \in \mathbb{k}$.
Case 3: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) \neq 0$. In this case this A_q-module cannot be extended to a D-module since we require $YX(v) = \tau(v)$.

In cases 1 and 2, we obtain an indecomposable D-module since it was indecomposable as an A_q-module to begin with.

Remark 6:
- *This module is irreducible as a D module if and only if $X \neq 0$ at any vector other than $(\mathfrak{m}_1, e_n)$ and Y, Y_1 are not simultaneously zero other than at $(\mathfrak{m}_0, e_0)$. If in addition, if $Y(v) = 0$ and $Y_1(w) = 0$ for some $v \neq w$ then we have an irreducible D-module which is not irreducible as an A_1 or an A_q-module.*
- *Another point to note, as we have seen in Case 3 above, not every A_q-module has a D-module structure.*
- *In Case 2 above, we could have $Y(\alpha(v)) = dw$ for some other w with $X(w) = 0, \tau(w) = 0$ thereby giving us yet another extension of an A_q-module to a D-module.*

3.1.6. *The module $V_q(\omega, w, f)$.* This family of A_q-decomposables arise when $|\omega| < \infty$ and $B_\omega \neq \emptyset$ and this is Family 2 as described in section 2.1. Let w be a word of length n where n is a multiple of m and let $f \in \mathbb{k}^*$. Consider n elements e_k where $k = 1, 2, \ldots, n$. For $\mathfrak{m} \in \omega$, let $V_{\mathfrak{m}}$ be a $\mathbb{k}_{\mathfrak{m}}$ vector space with basis $\{e_k \mid k \equiv j(mod\ m)\}$. The vector space $V_q(\omega, w, f) = \oplus_{\mathfrak{m} \in \omega} V_{\mathfrak{m}}$ has an A-module structure given by

$$X(\mathfrak{m}, e_k) = \begin{cases} t_{\mathfrak{m}}(\alpha(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \textit{ is not a break;} \\ (\alpha(\mathfrak{m}), e_{k+1}) & \textit{if } \mathfrak{m} \in B_\omega, k \neq n, z_{k+1} = x; \\ f(\alpha(\mathfrak{m}), e_1) & \textit{if } \mathfrak{m} \in B_\omega, k = n, z_1 = x; \\ 0 & \textit{otherwise;} \end{cases}$$

$$Y_1(\mathfrak{m}, e_k) = \begin{cases} (\alpha^{-1}(\mathfrak{m}), e_k) & \text{if } \alpha^{-1}(\mathfrak{m}) \text{ is not a break;} \\ (\alpha^{-1}(\mathfrak{m}), e_{k-1}) & \text{if } \alpha^{-1}((\mathfrak{m}) \in B_\omega, k \neq 1, z_k = y; \\ f(\alpha^{-1}(\mathfrak{m}), e_n) & \text{if } \alpha^{-1}(\mathfrak{m}) \in B, k = 1, z_1 = y, s = 1; \\ 0 & \text{otherwise;} \end{cases}$$

The module $V_q(\omega, w, f)$ is irreducible if $w = x^m$ or $w = y^m$. The extension of this action to an action of D follows the same pattern as in the previous case $V_q(\omega, j, w)$, section 3.1.5. That is,
Case 1: $X(v) = c\alpha(v) \neq 0$. Then, set $Y(\alpha(v)) = \frac{\tau(v)}{c}$.
Case 2: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) = 0$. In this case set $Y(\alpha(v)) = dv$ for any $d \in \Bbbk$.
Case 3: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y_1(\alpha(v)) = v$ and $\tau(v) \neq 0$. In this case this A_q-module cannot be extended to a D-module since we require $YX(v) = \tau(v)$. The extended D-module is also indecomposable.

Remark 7:
- *This module is irreducible as a D module if and only if $X = 0$ at at most one basis vector and Y, Y_1 are not simultaneously zero.*
- *Not every A_q-module has a D-module structure.*
- *In Case 2 above, we could have $Y(\alpha(v)) = dw$ for some other w with $X(w) = 0, \tau(w) = 0$ thereby giving us yet another extension of an A_q-module to a D-module.*

3.2. The list of A_1-indecomposables.

Now we go through the list of GWA-indecomposables as described in [2] and understand them as A_1-weight modules.

3.2.1. *The module $V_1(\omega, a)$.* Heret $|\omega| = \infty$ and $B_\omega = \emptyset$. Hence, $a \in \Bbbk \setminus \mathbb{Z}$ if the characteristic of $\Bbbk$ is 0, and $a \in \Bbbk \setminus \mathbb{Z}_p$ if the characteristic of $\Bbbk$ is $p > 0$. Here, $V_1(\omega, a) = \oplus_{i\in\mathbb{Z}} \Bbbk v_i$ where $\{v_i\}_{i\in\mathbb{Z}}$ is a basis, and the action is given by

$$X(v_i) = (a+i)v_{i+1}, \quad Y(v_i) = v_{i-1}, \quad f(\tau, \sigma)(v_i) = f(a+i, q^i b)v_i, \quad \text{for } f \in \Bbbk[\tau, \sigma, \sigma^{-1}].$$

For any $b \in \Bbbk^*$, we may extend the action of A_1 on $V_1(\omega, a)$ to an action of D by setting $Y_1(v_i) = \dfrac{(q^i b - 1)}{(a + i - 1)} v_{i-1}$. The resulting module is denoted $V_1(\omega, a, b)$ and is an irreducible D-module. By a suitable change of basis the picture for $V_1(\omega, a, b)$ is identical to the picture of $V_q(\omega, b, a)$ drawn in case 3.1.1.

3.2.2. *The module* $V_1(\omega, J, J')$ *when characteristic of* $\Bbbk$ *is* 0. Suppose that $|\omega| = \infty$ and $B_\omega \neq \emptyset$. That is, $\omega = \{\alpha^i(\tau - a, \sigma - b)\}_{i\in\mathbb{Z}}$. Since $B = \{(\tau, \sigma - c) \mid c \in \Bbbk\}$, we have $B_\omega \neq \emptyset$ if and only if $a \in \mathbb{Z}$. Without loss of generality, $a = 0$. That is, $\omega = \{\mathfrak{m}_i = (\tau - i, \sigma - q^i b)\}_{i\in\mathbb{Z}}$, and $B_\omega = \{\mathfrak{m}_0 = (\tau, \sigma - b)\}$. Set $B'_\omega = \{\mathfrak{m}_0, \mathfrak{m}_1\}$. Thus, we have the following options:

(1) $J = B'_\omega, J' = \{\mathfrak{m}_0\}, \mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \infty$

In this case, we have $V_1(\omega, J, J') = \oplus_{i=-\infty}^{\infty} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \in \mathbb{Z}$.

$$X(v_i) = \begin{cases} v_1 & \text{if } i = 0 \\ i v_{i+1} & \text{if } i \neq 0; \end{cases} \quad Y(v_i) = \begin{cases} 0 & \text{if } i = 1 \\ v_{i-1} & \text{if } i \neq 1 \end{cases};$$

$$\sigma(v_i) = q^i b v_i, \quad \tau(v_i) = i v_i \quad \forall i \in \mathbb{Z}.$$

Using the fact that $Y_1 X = q\sigma - 1$, for a fixed $b \in \Bbbk^*$ we define an action of Y_1 on $V_1(\omega, J, J')$ as follows: $Y_1(v_{i+1}) = \begin{cases} \dfrac{(q^{1+i}b - 1)}{i} v_i & \forall i \neq 0 \\ (qb - 1)v_0. & i = 0. \end{cases}$ The resulting D module, denoted by $V_1(\omega, J, J', b)$, is irreducible if $b \neq \frac{1}{q}$. If $b = q^i$ for some $i \in \mathbb{Z}, i \neq -1$, then this module is irreducible as a D module even though it is not irreducible as an A_1 module or an A_q module. The picture for this module is identical to the one drawn in subsection 3.1.1 for $a = 0$.

(2) $J = B'_\omega, J' = \emptyset, \mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \infty$.

In this case again, we have $V_1(\omega, J, J') = \oplus_{i=-\infty}^{\infty} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \in \mathbb{Z}$. Here,

$$X(v_i) = i v_{i+1}; \quad Y(v_i) = v_{i-1}; \quad \sigma(v_i) = q^i b v_i, \quad \tau(v_i) = i v_i \quad \forall i \in \mathbb{Z}.$$

Using the fact that $Y_1 X = q\sigma - 1$, we now define action of Y_1 on $V(\omega, J, J')$ by

$$Y_1(v_i) = \begin{cases} c v_0 & \text{if } i = 1, \text{ for some } c \in \Bbbk, \\ \left(\dfrac{q^i b - 1}{i - 1}\right) v_{i-1} & \text{if } i \neq 1. \end{cases}$$

Requiring that $XY_1(v_1) = (\sigma - 1)(v_1)$ gives us $b = q^{-1}$. Thus, we have the following:

$$\sigma(v_i) = q^{i-1} v_i; \quad Y_1(v_i) = \begin{cases} c v_0 & \text{if } i = 1, \text{ for some } c \in \Bbbk, \\ \left(\frac{q^{i-1} - 1}{i-1}\right) v_{i-1} & \text{if } i \neq 1. \end{cases}$$

For any fixed $c \in \Bbbk$, we have a D-module denoted by $V_1(\omega, J, J', c)$, which is indecomposable, but not irreducible.

Remark 8: *If $b \neq q^{-1}$, then $V_1(\omega, J, J')$ does not have a D-module structure.*

We may generalize this construction to the following D-module denoted by $V_1(\omega, J, J', c, d)$ for any $c, d \in \Bbbk$. Set $V_1(\omega, J, J', c, d) = \oplus_{i=-\infty}^{\infty} \Bbbk v_i$. Here,

$$X(v_i) = i v_{i+1}; \quad \sigma(v_i) = q^{i-1} v_i; \quad \tau(v_i) = i v_i \quad \forall i \in \mathbb{Z}.$$

$$Y(v_i) = \begin{cases} c v_0 & \text{if } i = 1, \text{ for some } c \in \Bbbk, \\ v_{i-1} & \text{if } i \neq 1; \end{cases}$$

$$Y_1(v_i) = \begin{cases} d v_0 & \text{if } i = 1, \text{ for some } d \in \Bbbk, \\ \left(\dfrac{q^{i-1} - 1}{i - 1}\right) v_{i-1} & \text{if } i \neq 1. \end{cases}$$

The D-module $V_1(\omega, J, J', c, d)$ is indecomposable. Its picture is identical to that of 3.1.2, case (2).

(3) $J = \{\mathfrak{m}_0\}, J' = \emptyset, \mathfrak{n}_0 = -\infty, \mathfrak{n}_1 = \mathfrak{m}_0$

In this case again, we have $V_1(\omega, J, J') = \oplus_{i \le 0} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \le 0$. Here, $Y(v_i) = v_{i-1}; \quad \sigma(v_i) = q^i b v_i, \quad \tau(v_i) = i v_i$ and

$$X(v_i) = \begin{cases} i v_{i+1} & \text{for } i \le -1 \\ 0 & \text{for } i = 0 \end{cases} \qquad \forall i \le 0.$$

Since $\tau - 1 \neq 0$, we can define Y_1 on on $V_1(\omega, J, J')$ by letting $Y_1 = Y\left(\dfrac{\sigma - 1}{\tau - 1}\right)$. This gives a module structure of D on on $V_1(\omega, J, J')$, and it is irreducible. The picture here is identical to that of 3.1.2, case (3).

(4) $J = \{\mathfrak{m}_1\}, J' = \emptyset, \mathfrak{n}_0 = \mathfrak{m}_0, \mathfrak{n}_1 = \infty$ In this case we have $V_1(\omega, J, J') = \oplus_{i \ge 1} \Bbbk v_i$. Set $\Bbbk v_i = K_{\mathfrak{m}_i}$ for $i \ge 1$. Here, $\sigma(v_i) = q^{i-1} b v_i, \quad \tau(v_i) = i v_i$ and

$$X(v_i) = i v_{i+1} \quad Y(v_i) = \begin{cases} v_{i-1} & \text{for } i \ge 2, \\ 0 & \text{for } i = 1 \end{cases}; \quad \forall i \ge 1.$$

Since $\tau - 1 \neq 0$ on $v_i, i \ge 2$, we can define Y_1 on on $V_1(\omega, J, J')$ by letting $Y_1 = Y\left(\dfrac{\sigma - 1}{\tau - 1}\right)$ on $v_i, i \ge 2$ and $Y_1(v_1) = 0$. This gives a module structure of D on on $V_1(\omega, J, J')$, and it is irreducible. Here the picture is identical to that of 3.1.2, case (4).

(5) $J = \emptyset, J' = \emptyset, \mathfrak{n}_0 = \mathfrak{m}_0, \mathfrak{n}_1 = \infty$. The resulting module is the same as the one above in case (4).

(6) $J=\emptyset, J'=\emptyset$, $\mathfrak{n}_0=-\infty, \mathfrak{n}_1=\infty$. The resulting module is the same as the one above in case (2).

3.2.3. *The module* $V_1(\omega, J, J')$ *when characteristic of* $\mathbb{k}$ *is nonzero.* Note that since $|\omega|=\infty$ and characteristic of $\mathbb{k}$ is $p>0$, this case arises when q is not a root of 1. Since D contains $A_1^q(\mathbb{k})$, we can extend only those A_1-modules which are infinite dimensional. Thus, we consider only those $J\subset B'_\omega$ for which $\mathfrak{n}_0=-\infty$ or $\mathfrak{n}_1=\infty$. We also have $a\in\mathbb{Z}_p$, $|B_\omega|=\infty$ and $B'_\omega=B_\omega$. Set $J'\subset J$ not containing the maximal element of J and we consider the indecomposable A_1-module $V_1(\omega, J, J')$. We point out that $V_1(\omega, J, J')$ is not irreducible A_1-module for any choice of J'.

Not every $V_q(\omega, J, J')$ can be extended to a D module. Using the condition $YX=\tau$ we see the following cases for any $v\neq 0, v\in V_{\mathfrak{m}}$:
Case 1: $X(v)=c\alpha(v)\neq 0$. Then, set $Y_1(\alpha(v))=\frac{(q\sigma-1)(v)}{c}$.
Case 2: For $\mathfrak{m}\in J\setminus J'$, $X(v)=0$, $Y(\alpha(v))=v$ and $q\sigma(v)=v$. In this case set $Y_1(\alpha(v))=dv$ for any $d\in\mathbb{k}$.
Case 3: For $\mathfrak{m}\in J\setminus J'$, $X(v)=0$, $Y(\alpha(v))=v$ and $q\sigma(v)\neq v$. In this case this A_1-module cannot be extended to a D-module since we require $Y_1X(v)=(q\sigma-1)(v)$. In particular, since q is not a root of 1, we have that $q\sigma=1$ can happen for at most one weight $\mathfrak{m}$. Thus, if $\mathfrak{m}_1,\mathfrak{m}_2\in J\setminus J'$ are such that $X(v_1)=X(v_2)=0$ for $v_1,v_2\neq 0, v_1\in\mathfrak{m}_1, v_2\in\mathfrak{m}_1$, then this A_1-module cannot be extended to a D-module.

Remark 9: *In case there is a D-module structure, then that structure is indecomposable D-module since it was indecomposable as an* A_1*-module to begin with. This module is irreducible as a D module if and only if* $X\neq 0$ *for any* $\mathfrak{m}\neq\mathfrak{n}_1$ *and* Y, Y_1 *are not simultaneously zero unless* $\alpha^{-1}(\mathfrak{m})=\mathfrak{n}_0$*. Further, if* $X\neq 0$ *for any* $\mathfrak{m}\neq\mathfrak{n}_1$ *and* $Y(v)=Y_1(w)=0$ *for nonzero* $v,w,v\neq w,v,w\notin V_{\alpha(\mathfrak{n}_0)}$ *then we have an irreducible D-module which is not irreducible as an* A_1 *or an* A_q*-module.*

3.2.4. *The module* $V_1(\omega, f, a)$ *for* $a\notin\mathbb{Z}_p$*.* This family of A_q-decomposables arise when $|\omega|<\infty$ and $B_\omega=\emptyset$. In particular, the characteristic of $\mathbb{k}=p>0$, and q is a root of 1 Since $B_\omega=\emptyset$, we have $\omega=\alpha^i(\mathfrak{m})$ where $\mathfrak{m}=(\tau-a,\sigma-b)$ where $a,b\in\mathbb{k}, b\neq 0, a\notin\mathbb{Z}_p$. Suppose $|\omega|=r$ (therefore, $q^r=1$ and p/r). For any $f\in\mathbb{k}^*$, let $V_1(\omega, f, a)=\oplus_{1\leq i\leq r}\mathbb{k}v_i$ be an r-dimensional vector space and give it A_1-module structure as described in section 2.1, with $A=A_1$, $T=Y$, and

$V(\omega,f,a)=V(\omega,f)$ setting

$$\begin{aligned}\sigma(v_i)&=q^{i-1}bv_i;\\ \tau(v_i)&=(a+i-1)v_i;\end{aligned}\qquad X(v_i)=\begin{cases}(a+i-1)v_{i+1} & \textit{if } i<r,\\ f(a-1)v_1 & \textit{if } i=r;\end{cases}$$

$$Y(v_i)=\begin{cases}v_{i-1} & \textit{if } i>1,\\ \frac{1}{f}v_r & \textit{if } i=1.\end{cases}$$

Since action of X is an isomorphism, we extend the action of A_1 to an action of D by using $Y_1X=q\sigma-1$ and setting

$$Y_1(v_{i+1})=\begin{cases}\dfrac{q^ib-1}{a+i-1}v_i & \textit{if } i>1;\\ \dfrac{b-1}{f(a-1)}v_r & \textit{if } i=1.\end{cases}$$

We denote the resulting irreducible D-module by $V_1(\omega,f,a,b)$. With a change of basis we see the picture as given in section 3.1.4

3.2.5. *The module* $V_1(\omega,j,w)$. This family of A_1-decomposables arise when $|\omega|<\infty$ and $B_\omega\neq\emptyset$ and this is Family 1 as described in section 2.1. Let $|B_\omega|=m>0$ (that is, $a\in\mathbb{Z}_p$). The one-to-one correspondence $\mathbb{Z}_m\to B_m$ and $j(\mathfrak{m})\in\mathbb{Z}_m$ are as described in section 2.1. Let x,y be two noncommuting variables.

Fix $j\in\mathbb{Z}_m$ and $w=z_1z_2\cdots z_n$ be a word of length $n\geq 1$ where each $z_i\in\{x,y\}$. Let $e_0,e_1,\dots,e_n$ be $n+1$ symbols. For each $\mathfrak{m}\in\omega$, let $V_{\mathfrak{m}}$ be a vector space over $\Bbbk_{\mathfrak{m}}$ with basis $\{(\mathfrak{m},e_k)\mid k+j=j(\mathfrak{m})\in\mathbb{Z}_m\}$. Put $V_1(\omega,j,w)=\oplus_{\mathfrak{m}\in\omega}V_{\mathfrak{m}}$ and it has an A_1-module structure given by

$$X(\mathfrak{m},e_k)=\begin{cases}t_{\mathfrak{m}}(\alpha(\mathfrak{m}),e_k) & \textit{if } \mathfrak{m}\notin B,\\ (\alpha(\mathfrak{m}),e_{k+1}) & \textit{if } \mathfrak{m}\in B_\omega \textit{ and } z_{k+1}=x,\\ 0 & \textit{otherwise};\end{cases}$$

$$Y(\mathfrak{m},e_k)=\begin{cases}(\alpha^{-1}(\mathfrak{m}),e_k) & \textit{if } \mathfrak{m}\notin B,\\ (\alpha^{-1}(\mathfrak{m}),e_{k-1}) & \textit{if } \mathfrak{m}\in B_\omega \textit{ and } z_k=y,\\ 0 & \textit{otherwise.}\end{cases}$$

The module $V_1(\omega,j,w)$ is irreducible as an A_1-module when w is the empty word. Not every $V_1(\omega,j,w)$ can be extended to a D module. Using the condition $Y_1X=q\sigma-1$ we see the following cases for any $v\neq 0, v\in V_{\mathfrak{m}}$:

Case 1: $X(v) = c\alpha(v) \neq 0$. Then, set $Y_1(\alpha(v)) = \frac{(q\sigma-1)(v)}{c}$.
Case 2: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y(\alpha(v)) = v$ and $q\sigma(v) = v$. In this case set $Y_1(\alpha(v)) = dv$ for any $d \in \Bbbk$.
Case 3: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y(\alpha(v)) = v$ and $q\sigma(v) \neq v$. In this case this A_1-module cannot be extended to a D-module since we require $Y_1X(v) = (q\sigma - 1)(v)$.

In cases 1 and 2, we obtain an indecomposable D-module since it was indecomposable as an A_1-module to begin with.

Remark 10:

- *This module is irreducible as a D module if and only if $X \neq 0$ at any vector other than $(\mathfrak{m}_1, e_n)$ and Y, Y_1 are not simultaneously zero other than at $(\mathfrak{m}_0, e_0)$. If in addition, if $Y(v) = 0$ and $Y_1(w) = 0$ for some $v \neq w$ then we have an irreducible D-module which is not irreducible as an A_1 or an A_q-module.*
- *Another point to note, as we have seen in Case 3 above, not every A_1-module has a D-module structure.*
- *In Case 2 above, we could have $Y_1(\alpha(v)) = dw$ for some other w with $X(w) = 0, q\sigma(w) = w$ thereby giving us yet another extension of an A_1-module to a D-module.*

3.2.6. *The module $V_1(\omega, w, f)$.* This family of A_1-decomposables arise when $|\omega| < \infty$ and $B_\omega \neq \emptyset$ and this is Family 2 as described in section 2.1. Let w be a word of length n where n is a multiple of m and let $f \in \Bbbk^*$. Consider n elements e_k where $k = 1, 2, \ldots, n$. For $\mathfrak{m} \in \omega$, let $V_\mathfrak{m}$ be a $\Bbbk_\mathfrak{m}$ vector space with basis $\{e_k \mid k \equiv j(mod\ m)\}$. The vector space $V_1(\omega, w, f) = \oplus_{\mathfrak{m}\in\omega} V_\mathfrak{m}$ has an A-module structure given by

$$X(\mathfrak{m}, e_k) = \begin{cases} t_\mathfrak{m}(\alpha(\mathfrak{m}), e_k) & \textit{if } \mathfrak{m} \textit{ is not a break;} \\ (\alpha(\mathfrak{m}), e_{k+1}) & \textit{if } \mathfrak{m} \in B_\omega, k \neq n, z_{k+1} = x; \\ f(\alpha(\mathfrak{m}), e_1) & \textit{if } \mathfrak{m} \in B, k = n, z_1 = x; \\ 0 & \textit{otherwise;} \end{cases}$$

$$Y(\mathfrak{m}, e_k) = \begin{cases} (\alpha^{-1}(\mathfrak{m}), e_k) & \textit{if } \alpha^{-1}(\mathfrak{m}) \textit{ is not a break;} \\ (\alpha^{-1}(\mathfrak{m}), e_{k-1}) & \textit{if } \alpha^{-1}((\mathfrak{m}) \in B, k \neq 1, z_k = y; \\ f(\alpha(\mathfrak{m}), e_n) & \textit{if } \alpha^{-1}(\mathfrak{m}) \in B, k = 1, z_1 = y; \\ 0 & \textit{otherwise;} \end{cases}$$

The module $V_1(\omega, w, f)$ is irreducible if $w = x^m$ or $w = y^m$. The extension of this action to an action of D follows the same pattern as in the previous case $V_1(\omega, j, w)$, section 3.2.5. That is,
Case 1: $X(v) = c\alpha(v) \neq 0$. Then, set $Y_1(\alpha(v)) = \frac{(q\sigma-1)(v)}{c}$.
Case 2: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y(\alpha(v)) = v$ and $q\sigma(v) = v$. In this case set $Y_1(\alpha(v)) = dv$ for any $d \in \Bbbk$.
Case 3: For $\mathfrak{m} \in B_\omega$, $X(v) = 0$, $Y(\alpha(v)) = v$ and $q\sigma(v) \neq v$. In this case this A_1-module cannot be extended to a D-module since we require $Y_1X(v) = (q\sigma - 1)(v)$.

When one can extend to a D-structure, the extended D-module is also indecomposable.

Remark 11:

- *This module is irreducible as a D module if and only if $X = 0$ at at most one basis vector and Y, Y_1 are not simultaneously zero.*
- *Not every A_1-module has a D-module structure.*
- *In Case 2 above, we could have $Y_1(\alpha(v)) = dw$ for some other w with $X(w) = 0, q\sigma(w) = w$ thereby giving us yet another extension of an A_1-module to a D-module.*

4. Irreducible D-modules

Proposition 1: *All nonzero weight-spaces of an irreducible weight D-module are* 1*-dimensional.*

Proof. Consider the following quiver Q



subject to all the relations inherited from D. Let $\Bbbk Q$ be the path algebra of Q. Fix an ith vertex and consider the cyclic subalgebra $Q(i,i)$ of $\Bbbk Q$ which consists of all paths that start and end in i. It is a standard fact that the restriction of any irreducible $\Bbbk Q$-module V onto $Q(i,i)$ is irreducible. But $Q(i,i) = \Bbbk[\sigma, \tau]$.

Since $\Bbbk$ is algebraically closed then any irreducible module over $Q(i,i)$ is 1-dimensional. ∎

Theorem 1: *Let V be an irreducible D weight module. Then V is indecomposable as an A_q as well as an A_1 weight module.*

Proof. Let $V = \oplus_{\mathfrak{m}\in\omega} V_{\mathfrak{m}}$ be an irreducible D weight module where ω is one orbit.

Case: X is injective on V. If ω is cyclic, then V is finite dimensional and every nonzero weight space is one dimensional. Let $v_0 \in \mathfrak{m}_0 \in \omega, v_0 \neq 0$. Then $X^i(v_0) \in \alpha^i(\mathfrak{m}_0)$ and $X^i(v_0) \neq 0$ for every $i \geq 0$. That is, if $\omega = \{\alpha^i(\mathfrak{m}_0) \mid 0 \leq i \leq r\}$, then V has basis $\{X^i(v_0) \mid 0 \leq i \leq r\}$, and the only X-invariant subspaces are $\{0\}$ and V. That is, V is indecomposable as an A_1 and as an A_q module.

If ω is linear and suppose there exists a $v_0 \in V, v_0 \neq 0$ such that the preimage $X^{-1}(v_0) = \emptyset$. If $Y(v_0) = w \neq 0$, then $XY(v_0) = (\tau - 1)(v_0)$. If $(\tau-1)(v_0) = cv_0, c \neq 0$, then $X(\frac{w}{c}) = v_0$ contradicting the fact that $X^{-1}(v_0) = \emptyset$. Thus, $XY(v_0) = (\tau - 1)(v_0) = 0$. If $w \neq 0$, then $X(w) = 0$ contradicting the injectivity of X. Thus, $Y(v_0) = 0$. Similar argument gives us $Y_1(v_0) = 0$. In other words, the span of $\{X^i(v_0) \mid i \geq 0\}$ is a D-submodule of V, and hence $V = \oplus_{i=0}^{\infty} \Bbbk X^i(v_0)$ and therefore indecomposable as A_1 and A_q module.

Suppose the preimage $X^{-1}(v) \neq \emptyset$ for any weight vector v. Let $v_0 \in V_{\mathfrak{m}_0}, v_0 \neq 0$ be a weight vector. Then $V_{\alpha^i(\mathfrak{m}_0)} = \Bbbk X^i(v_0)$ for any $i \in \mathbb{Z}$. Note that since X is injective, $X^i(v_0)$ is a singleton set for $i < 0$. Thus, V is indecomposable as A_1 and A_q module.

Case: $X(v_0) = 0$ for a nonzero vector $v_0 \in V_{\mathfrak{m}_0}$ for some $\mathfrak{m}_0 \in \omega$. Then $V = D \cdot v_0 = S \cdot v_0$ where S is the subalgebra of D generated by Y, Y_1 over R.

Suppose $|\omega| = \infty$. That is, $V = \oplus_{i \leq 0} V_{\alpha^i(\mathfrak{m}_0)} \neq 0$. Suppose $X : V_{\alpha^i(\mathfrak{m}_0)} \to V_{\alpha^{i+1}(\mathfrak{m}_0)}$ is zero on these one-dimensional spaces for some $i < 0$. Then, $W = \oplus_{j \leq i} V_{\alpha^j(\mathfrak{m}_0)}$ is a D-submodule of V. Therefore, $X : V_{\alpha^i(\mathfrak{m}_0)} \to V_{\alpha^{i+1}(\mathfrak{m}_0)}$ is an isomorphism for every $i < 0$. For any A_1 or A_q submodule W of V, if $v_i \in W$, then $X^j(v_i) \in W, \quad \forall j \geq 0$. In particular $v_0 \in W$. Thus V is A_1 and A_q indecomposable.

Suppose $|\omega| = r+1 < \infty$. Let $V = V_{\mathfrak{m}_r} \oplus V_{\mathfrak{m}_{r-1}} \oplus \ldots \oplus V_{\mathfrak{m}_0}$. Suppose $V_{\mathfrak{m}_i} = 0$ for some $i, 0 < i \leq r$, then $W = V_{\mathfrak{m}_{i-1}} \oplus V_{\mathfrak{m}_{i-2}} \oplus \ldots \oplus V_{\mathfrak{m}_0}$ is a D-submodule of V. Therefore, we may assume that every $V_{\mathfrak{m}_i}$ is one dimensional for $0 \leq i \leq r$ and let u_i be a basis vector for $V_{\mathfrak{m}_i}$ for each $0 \leq i \leq r$. Note that $Y, Y_1 : V_{\mathfrak{m}_r} \to V_{\mathfrak{m}_0}$

since $\alpha^{-1}(\mathfrak{m}_r) = \mathfrak{m}_0$. Since $XY(u_r) = XY_1(u_r) = 0$ and $XY = \tau - 1$ and $XY_1 = \sigma - 1$, we have $\mathfrak{m}_r = (\tau - 1, \sigma - 1)$. If $X : V_{\mathfrak{m}_i} \to V_{\mathfrak{m}_{i-1}}$ is the zero map for any $i, 0 < i \leq r$, then again $\mathfrak{m}_{i-1} = (\tau - 1, \sigma - 1)$ contradicting the fact that ω is a single orbit.

Therefore, $X : V_{\mathfrak{m}_i} \to V_{\mathfrak{m}_{i-1}}$ is an isomorphism for every $0 < i \leq r$. Then $V = \oplus_{i=0}^{r} \Bbbk X^i(u_r)$; in other words, every X-invariant subspace of V contains v_0, thereby making V an indecomposable A_1 and an A_q module. ∎

4.1. Characteristic of $\Bbbk$ is 0. Here $\mathfrak{m} = (\tau - a, \sigma - b)$ for some $a, b \in \Bbbk$, with $b \neq 0$. Note that $(q\sigma - 1) \in \mathfrak{m}$ if and only if $b = \dfrac{1}{q}$ and $\tau \in \mathfrak{m}$ if and only if $a = 0$. In other words, the set of breaks $B = \{(\tau - a, \sigma - \frac{1}{q}) \mid a \in \Bbbk\}$ when studying A_q weight modules, and $B = \{(\tau, \sigma - b) \mid b \in \Bbbk\}$ when studying A_1 weight modules.

Fix $\mathfrak{m} = (\tau - a, \sigma - b)$. Then, $\alpha^k(\mathfrak{m}) = (\tau - a - k, \frac{\sigma}{q^k} - b)$. That is, $\omega = \{(\tau - a - k, \frac{\sigma}{q^k} - b) \mid k \in \mathbb{Z}\}$. If $\Bbbk$ is of characteristic 0 or if q is not a root of unity, then $|\omega| = \infty$. Note, $(q\sigma - 1) \in \alpha^k(\mathfrak{m})$ if and only if $b = q^{-k-1}$ (respectively, $\tau \in \alpha^k(\mathfrak{m})$ if and only if $a = -k$) for $k \in \mathbb{Z}$. That is, $|B_\omega| \leq 1$ since $B_\omega = \emptyset$ if $b \notin \{q^i\}_{i\in\mathbb{Z}}$ or $B_\omega = (\tau - a - k, q\sigma - 1)$ if $b = q^{-k-1}$, while studying A_q-modules; or $B_\omega = \emptyset$ if $a \notin \mathbb{Z}$ or $B_\omega = (\tau, \sigma - q^k b)$ if $a = k$ while studying A_1-modules.

We now consider the following families:

Family I: Let V be an irreducible A_q weight module.

Case: q is a root of 1. Here, by Theorem (5.8) of [2] V is isomorphic to $V_q(\omega, b)$ for $b \in \Bbbk^* \setminus \{q^i\}_{i\in\mathbb{Z}}$. For any $a \in \Bbbk$, the module $V_q(\omega, b)$ can be given an irreducible D-weight module structure, and we denote this D module by $V_q(\omega, b, a)$. The details are given in section 3.1, case 3.1.1.

Case: q is not a root of 1. In this case we have more families of irreducible modules. Here, as an irreducibel A_q-module, V is one of the three: $V_q(\omega, b)$ for $b \in \Bbbk^* \setminus \{q^i\}_{i\in\mathbb{Z}}$ (described in section 3.1, case 3.1.1), or $V_q(\omega, J, J')$ (described in section 3.1 case 3.1.2) subcases (3) and (4). For any $a \in \Bbbk$, these A_q-modules are extended to irreducible D weight modules; these extensions, denoted $V_q(\omega, b, a), V_q(\omega, J, J', a)$, are also described in the mentioned sections.

Family II: Let V be an irreducible A_1 weight module. Here, V is one of the three: $V_1(\omega, b)$ (described in section 3.2.1) and $V_1(\omega, J, J', b)$ cases (3) and (4) (described in section 3.2.2 cases (3) and (4)). Since these modules are A_1 irreducible, they are D-irreducible whether or not q is a root of unity.

Family III: Irreducible D weight modules which do not arise from the families I and II above A D weight module is also an A_q and an A_1 weight module. Note that an irreducible D-weight module need not result in an irreducible A_q or an irreducible A_1 weight module. For instance, the module $V_q(\omega, J, J', a)$ described in 3.1.2 case (1) is an irreducible D weight module which does not arise from the two families I and II. But by Theorem 1, every irreducible D-module results in an indecomposable A_1 and an indecomposable A_q-module. So, we start with those modules V which are indecomposable (but not irreducible) as A_1-modules, and which can be extended to irreducible D-modules while ensuring that they stay indecomposable (but not irreducible) as A_q-modules.

Thus, $|\omega| = \infty$, and $B_\omega \neq \emptyset$, where $\omega = \{\alpha^i(\tau - a, \sigma - b)\}$. Here, refer to section 3.2.2. We see $B_\omega \neq \emptyset$ implies that $a \in \mathbb{Z}$. We obtain irreducible D-modules in two ways. One is described in case (1), for $b \in \Bbbk \setminus \{q^{a-1}\}$. The other, and its generalization, is described in case (2) for $b = q^{a-1}$. Note that in these cases, q may or may not be a root of unit (refer to 3.1.3).

Theorem 2: *When characteristic of $\Bbbk$ is 0 the irreducible D-modules are as described in families I, II, and III.*

Remark 12: *Not every indecomposable A_1-module can be given a D-module structure, (see Remark 8, section 3.2.2). Likewise, not every indecomposable A_q-module can be given a D-module structure, (see Remark 2, section 3.1.2, and Remark 5 , section 3.1.3).*

Remark 13: *The key point in Theorem 1 is Proposition 1; that is, every weight-space of an irreducible D weight module is one dimensional. Note that an indecomposable D weight module need not satisfy this property. For example, let $V(c,d) = (\oplus_{i\leq -1}\Bbbk u_i) \oplus (\oplus_{i\leq -1}\Bbbk w_i) \oplus (\oplus_{i\geq 0}\Bbbk v_i)$ be given a D-module structure as follows:*

$$\sigma(x) = \begin{cases} \frac{x}{q^i} & \text{if } x = u_{-i}, \\ \frac{x}{q^i} & \text{if } x = w_{-i}, \\ q^i x & \text{if } x = v_i; \end{cases} \quad X(x) = \begin{cases} u_{i+1} & \text{if } x = u_i, i \neq -1, \\ w_{i+1} & \text{if } x = w_i, i \neq -1, \\ 0 & \text{if } x = w_{-1} \text{ or } x = u_{-1}, \\ v_{i+1} & \text{if } x = v_i; \end{cases}$$

$$\tau(x) = \begin{cases} -ix & \textit{if } x = u_{-i}, \\ -ix & \textit{if } x = w_{-i}, \\ ix & \textit{if } x = v_i; \end{cases} \quad Y(x) = \begin{cases} (i-1)u_{i-1} & \textit{if } x = u_i, \\ (i-1)w_{i-1} & \textit{if } x = w_i, \\ (i-1)v_{i-1} & \textit{if } x = v_i, i \neq 0, \\ cu_{-1} & \textit{if } x = v_0; \end{cases}$$

$$Y_1(x) = \begin{cases} (q^{i-1}-1)u_{i-1} & \textit{if } x = u_i, \\ (q^{i-1}-1)w_{i-1} & \textit{if } x = w_i, \\ (q^{i-1}-1)v_{i-1} & \textit{if } x = v_i, i \neq 0, \\ dw_{-1} & \textit{if } x = v_0; \end{cases}$$

One can see that $V(c,d), c, d \neq 0$ is an indecomposable D-weight module, with weight-spaces $V_{\mathfrak{m}_i} = \Bbbk u_i \oplus \Bbbk w_i$ for $i \leq -1$ where $\mathfrak{m}_i$ is the maximal ideal $(\tau - i, \sigma - \frac{1}{q^i})$, and $V_{\mathfrak{m}_i} = \Bbbk v_i$ for $i \geq 0$. The picture for this is as follows:



4.2. Characteristic of $\Bbbk$ is nonzero and q is not a root of 1. Here again we go through three families: Family I of irreducible A_q modules, Family II of irreducible A_1 modules, and Family III of indecomposable A_q and A_1 modules which are not irreducible in either case.

Family I: Here, as an irreducible A_q-module, V is one of the three: $V_q(\omega, b)$ for $b \in \Bbbk^* \setminus \{q^i\}_{i \in \mathbb{Z}}$ (described in section 3.1, case 3.1.1), or $V_q(\omega, J, J')$ (described in section 3.1 case 3.1.2) subcases (3) and (4). For any $a \in \Bbbk$, these A_q-modules are extended to irreducible D weight modules; these extensions, denoted $V_q(\omega, b, a), V_q(\omega, J, J', a)$, are also described in the mentioned sections. When $a \in \Bbbk \setminus \mathbb{Z}_p$, then the resulting D-modules are also A_1-modules. If $a \in \mathbb{Z}_p$ then the resulting modules are indecomposable A_1-modules.

Family II: Let V be an irreducible A_1 weight module. Here, V is one of the three: $V_1(\omega, b, a)$ (described in section 3.2.1) and $V_1(\omega, J, J')$ (described in section 3.2.3). These are D-irreducible weight modules.

Family III: Irreducible D weight modules which do not arise from the families I and II above Here we start with indecomposable A_q-modules and extend them to irreducible D-modules while keeping A_1-structure indecomposable, although not irreducible. These appear in two situations of section 3.1.2: One is described in case (1) with $a \neq 0$; the other is descibed in case (2) and its generalization with $c = 0, d \neq 0$.

Theorem 3: *When characteristic of* $\Bbbk$ *is not zero, and* q *is not a root of* 1 *the irreducible* D*-modules are as described in families I, II, and III.*

4.3. Characteristic of $\Bbbk$ is nonzero and q is a root of 1. In this case, we see that $|\omega| < \infty$. Thus, we may describe families I and II as before.

Family I: When $B_\omega = \emptyset$, we get the module $V_q(\omega, f, b)$ for $b \notin \{q^i\}_{i \in \mathbb{Z}}$ and $f \in \Bbbk^*$ described in section 3.1.4. This is an irreducible A_q-module and extends to an irreducible D-module. When $B_\omega \neq \emptyset$, we have two families: The first family is the module $V_q(\omega, j, w)$ which is irreducible for any $j \in \mathbb{Z}_m$ and w the empty word; this module is described described in section 3.1.5. While not every $V_q(\omega, j, w)$ can be extended to a D-module, when w is the empty word, this module can be extended to a D-module which will be irreducible. The second family is the module $V_q(\omega, w, f)$ which is irreducible for the word $w = x^m$ or $w = y^m$; this module is described in section 3.1.6. When $w = x^m$, this module can be extended to a D-module. But if $w = y^m$, then this module may or may not be extended to a D-module.

Family II: As in the preceding paragraph, we have three families of irreducible A_1 modules which may be extended to an irreducible D-module. The module $V_1(\omega, f, a)$ for $a \notin \mathbb{Z}_p$ is described in section 3.2.4 and arises when $B_\omega = \emptyset$. This is an irreducible A_1 module, and can be extended to a D-module.

The module $V_1(\omega, j, w)$ is described in section 3.2.5 and arises when $B_\omega \neq \emptyset$. This is an irreducible A_1-module if and only if w is the empty word, and it can then be extended to a D-module.

The module $V_1(\omega, w, f)$ is described in section 3.2.6 and arises when $B_\omega \neq \emptyset$. This is an irreducible A_1 module if and only if $w = x^m$ or $x = y^m$. When $w = x^m$, this module can be extended to an irreducible D-module. When $x = y^m$, this module may or may not be extended to a D-module.

Family III: We start with indecomposable A_1-modules (but not irreducible) and extend in such a fashion that the resultant modules are indecomposable A_q-modules (but not irreducible) and are irreducible as D modules. Thus, we look at the case where $|\omega| < \infty$ and $B_\omega \neq \emptyset$. As has been explained in the proof of Theorem 1 the kernel of X is one-dimensional. By Proposition 1 every weight space is also one-dimensional. In other words, when we consider module $V_q(\omega, j, w)$ (section 3.1.5) or $V_1(\omega, j, w)$ (section 3.2.5) we need to consider $w = x^m$ where $m = |B_\omega|$. Both these families may or may not be extended to families of D-modules. If extended, the extended families are irreducible if and only if Y and Y_1 are not simultaneously zero except at $(\mathfrak{m}_0, e_0)$.

Similarly, we consider those modules $V_q(\omega, w, f)$ (section 3.2.6) and $V_1(\omega, w, f)$ (section 3.2.6). These modules are irreducible as A_q, and respectively A_1 modules if and only if $w = x^m$ or $w = y^m$. Since we require the dimension of the kernel of X to be at most 1, and the dimension of every weight space be equal to 1, we are forced to require that w be a word of degree m in variables x, y with degree in y being exactly 1. Both these families may or may not be extended to families of D-modules. If extended, the extended families are irreducible if and only if Y and Y_1 are not simultaneously zero except at the same edge as the edge $X = 0$.

THEOREM 4: *When characteristic of $\Bbbk$ is nonzero and q is a root of 1, the irreducible D weight modules are those described in families I ,II, and III.*

4.4. A FAMILY OF INDECOMPOSABLE D-MODULES WHICH ARE DECOMPOSABLE AS A_q AND A_1 MODULES. Throughout this section we assume that the characteristic of $\Bbbk$ is $p > 0$ and q is a root of 1 of order p.

In this section we present a family of finite dimensional indecomposable D-modules which are decomposable as A_q and as A_1 modules. First, we present an example.

4.4.1. *Characteristic of $\Bbbk$ is 3, and $q^3 = 1$.* Let $\mathfrak{m}_0 = (\tau - 1, \sigma - 1)$. Then $\alpha^i(\mathfrak{m}_0) = \{\mathfrak{m}_0, \mathfrak{m}_1, \mathfrak{m}_2\}$ with $\mathfrak{m}_i = (\tau - i - 1, \sigma - q^i)$. Fix $a, b, c, d \in \Bbbk^*$. Let $V = \oplus_{i=0}^{2} V_i$ where each V_i is a four dimensional $\Bbbk_{m_i}$ module with basis $\{(i, e_1), (i, e_2), (i, e_3), (i, e_4)\}$. This D module can be visualized as follows:



That is, the action of D on V is defined as follows:

$$X(i, e_j) = \begin{cases} (i+1, e_j) & \textit{if } i \neq 2, \\ 0 & \textit{if } i = 2; \end{cases}$$

$$Y(i, e_j) = \begin{cases} i(i-1, e_j) & \textit{if } i \neq 0, \\ a(2, e_1) & \textit{if } i = 0, j = 1, \\ (2, e_3) & \textit{if } i = 0, j = 2, \\ c(2, e_3) & \textit{if } i = 0, j = 3, \\ (2, e_1) & \textit{if } i = 0, j = 4; \end{cases}$$

$$Y_1(i,e_j)=\begin{cases}(q^i-1)(i-1,e_j) & \textit{if } i\neq 0,\\ (2,e_2) & \textit{if } i=0, j=1,\\ b(2,e_2) & \textit{if } i=0, j=2,\\ (2,e_4) & \textit{if } i=0, j=3,\\ d(2,e_4) & \textit{if } i=0, j=4.\end{cases}$$

While V is an indecomposable D-module, as an A_1 module V decomposes as $V=(W_1\oplus W_4)\oplus(W_2\oplus W_3)$ and as an A_q-module V decomposes as $V=(W_1\oplus W_2)\oplus(W_3\oplus W_4)$ where each W_i is the three dimensional vector space spanned by $\{(0,e_i),(1,e_i),(2,e_i)\}$ for $i=0,\dots,4$.

4.4.2. *Generalization.* The above example 4.4.1 can be generalized. Let $m\geq 4$ be an even natural number and $a_1,a_2,\dots,a_m\in\Bbbk^*$. Let $\mathfrak{m}_0=(\tau-1,\sigma-1)$, and $\mathfrak{m}_i=\alpha^i(\mathfrak{m}_0)$ for $0\leq i\leq p-1$. For each $0\leq i\leq p-1$ let V_i be the $\Bbbk_{\mathfrak{m}_i}$ with basis (i,e_j) for $1\leq j\leq m$. Let $V(m;a_1,a_2,\dots,a_m)=\oplus_{i=0}^{p-1}V_i$ and give it D-action by $X(i,e_j)=\begin{cases}(i+1,e_j) & \textit{if } i\neq p-1,\\ 0 & \textit{if } i=p-1;\end{cases}$

$$Y(i,e_j)=\begin{cases}i(i-1,e_j) & \textit{if } i\neq 0,\\ a_{2k-1}(p-1,e_j) & \textit{if } i=0, j=2k-1,\\ (p-1,e_{j+1}) & \textit{if } i=0, j=2k<m,\\ (p-1,e_1) & \textit{if } i=0, j=m;\end{cases}$$

$$Y_1(i,e_j)=\begin{cases}(q^i-1)(i-1,e_j) & \textit{if } i\neq 0,\\ a_{2k}(p-1,e_j) & \textit{if } i=0, j=2k,\\ (p-1,e_{j+1}) & \textit{if } i=0, j=2k-1,\end{cases}$$

where k is a natural number, $1\leq k\leq\frac{m}{2}$. This action results in an indecomposable D-module structure on $V(m;a_1,a_2,\dots,a_m)$. For each $1\leq j\leq m$, let W_j be the vector space with basis $\{(i,e_j)\mid 0\leq i\leq p-1\}$.

Note, for each $k, 1\leq k\leq\frac{m}{2}$, the vector space $W_{2k-1}\oplus W_{2k}$ is an A_q module, and viewed as an A_q-module, $V(m;a_1,a_2,\dots,a_m)$ is decomposed into a direct sum of A_q-submodules:

$$V(m;a_1,a_2,\dots,a_m)=\oplus_{k=1}^{\frac{m}{2}}(W_{2k-1}\oplus W_{2k}).$$

Similarly, for each $k, 1 \leq k < \frac{m}{2}$, the vector space $W_{2k} \oplus W_{2k+1}$ is an A_1 module; in addition, the vector space $W_m \oplus W_1$ is an A_1-module. Further, $V(m; a_1, a_2, \dots, a_m)$ is decomposed into a direct sum of A_1-submodules:

$$V(m; a_1, a_2, \dots, a_m) = \oplus_{k=1}^{\frac{m}{2}-1}(W_{2k} \oplus W_{2k+1}) \oplus (W_m \oplus W_1).$$

We thus have,

Example 1: For any even natural number $m \geq 4$, and $a_1, a_2, \dots, a_m \in \Bbbk^*$, there is an indecomposable D-weight module $V(m; a_1, a_2, \dots, a_m)$ which is decomposable as an A_q as well as an A_1 weight module into $\dfrac{m}{2}$ components.

This example leads us to the contruction of a family of indecomposable D modules as follows: Let Y_1, Y be two noncommuting variables, $z \in \{Y_1, Y\}$ and $a \in \Bbbk^*$. As above, let $\mathfrak{m}_0 = (\tau - 1, \sigma - 1)$, and $\mathfrak{m}_i = \alpha^i(\mathfrak{m}_0)$ for $0 \leq i \leq p-1$. Let $W^{z,a} = \oplus_{i=0}^{p-1}\Bbbk v_i$ where v_i is of weight $\mathfrak{m}_i$. Then $W^{z,a}$ has a D-module structure given by $X(v_i) = \begin{cases} v_{i+1} & \textit{if } i < p-1, \\ 0 & \textit{if } i = p-1; \end{cases}$

$$Y(v_i) = \begin{cases} iv_{i-1} & \textit{if } i > 0, \\ av_{p-1} & \textit{if } i = 0, \ \textit{and } z = Y, \\ 0 & \textit{if } i = 0, \ \textit{and } z = Y_1; \end{cases}$$

$$Y_1(v_i) = \begin{cases} (q^i - 1)v_{i-1} & \textit{if } i > 0, \\ av_{p-1} & \textit{if } i = 0, \ \textit{and } z = Y_1, \\ 0 & \textit{if } i = 0, \ \textit{and } z = Y. \end{cases}$$

Fix m a natural number, $a_1, a_2, \dots, a_m \in \Bbbk^*$, and $w = z_1 z_2 \cdots z_m$ a word of length m where $z_i \in \{Y_1, Y\}$. Let $V(m; w, a_1, \dots, a_m) = \oplus_{i=1}^{m} W^{z_i, a_i}$ as a vector space. We now connect each component to give $V(m; w, a_1, \dots, a_m)$ an indecomposable D module structure. Namely, $V(m; w, a_1, \dots, a_m)$ receives action of X from its action on each component W^{z_i,a_i}. The action of Y on W^{z_i,a_i} is also inherited if $z_i = Y$. If $z_i = Y_1$, then for $i > 0$, action of Y on $v_i \in W^{z_i,a_i}$ is inherited; for $v_0 \in W^{z_i,a_i}$ set $Y(v_0) = v_{p-1} \in W^{z_{i+1},a_{i+1}}$ when $1 \leq i < m$, and set $Y(v_0) = v_{p-1} \in W^{z_1,a_1}$ when $i = m$. Similarly, the action of Y_1 on W^{z_i,a_i} is also inherited if $z_i = Y_1$. If $z_i = Y$, then for $i > 0$, action of Y_1 on $v_i \in W^{z_i,a_i}$ is inherited; for $v_0 \in W^{z_i,a_i}$ set $Y_1(v_0) = v_{p-1} \in W^{z_{i+1},a_{i+1}}$ when $1 \leq i < m$, and set $Y_1(v_0) = v_{p-1} \in W^{z_1,a_1}$ when $i = m$.

For $1 \leq i \leq m$ if we view each W^{z_i,a_i} as a vertex, then $V(m; w, a_1, \ldots, a_m)$ is to be viewed as a directed cycle on m vertices with an edge Y (if $z_i = Y_1$) or Y_1 (if $z_i = Y$) joining vertex W^{z_i,a_i} to $W^{z_{i+1},a_{i+1}}$ if $i < m$ and joining W^{z_m,a_m} to W^{z_1,a_1} for $i = m$. As an examle, for word $w = Y_1YY_1$, and $a_1, a_2, a_3 \in \Bbbk^*$ the cycle would be



Here, every W^{z_i,a_i} is a p-dimensional vector space.

Theorem 5: *Let p denote the characteristic of $\Bbbk$ and assume that p is equal to the order of q as a root of 1. For natural number m, $a_1, a_2, \ldots, a_m \in \Bbbk^*$, and any word $w = z_1z_2\cdots z_m$ with $z_i \in \{Y_1, Y\}$ there is an irreducible D-weight module $V(m; w, a_1, a_2, \ldots, a_m)$ of dimension pm. We also have:*

- ***Case** $m = 1$**:** Here $V(m; w, a_1, a_2, \ldots, a_m)$ is an irreducible A_1 and A_q module.*
- ***Case** $m = 1, 2, 3$**:** Here $V(m; w, a_1, a_2, \ldots, a_m)$ is indecomposable as an A_1 or an A_q module.*
- ***Case** $m \geq 4$**:** If $w = Y_1^m$, then $V(m; w, a_1, a_2, \ldots, a_m)$ is indecomposable as an A_1 module, but decomposable as an A_q module. Similarly, if $w = Y^m$, then $V(m; w, a_1, a_2, \ldots, a_m)$ is indecomposable as an A_q module, but decomposable as an A_1 module.*
- ***Case** $m \geq 4$**:** If the degree of w in Y_1 is 1 then $V(m; w, a_1, a_2, \ldots, a_m)$ is indecomposable as an A_q module, but decomposable as an A_1 module.Similarly, if the degree of w in Y is 1 then $V(m; w, a_1, a_2, \ldots, a_m)$ is indecomposable as an A_1 module, but decomposable as an A_q module.*
- ***Case** $m \geq 4$**:** If w is of degree at least 2 in y and at least degree 2 in Y_1, then $V(m; w, a_1, a_2, \ldots, a_m)$ is decomposable as an A_1 and an A_q module.*

While we do not have any general statements of indecomposable finite dimensional D-modules we can say the following:

Theorem 6: *Let $V = \oplus_{\mathfrak{m}\in\omega} V_{\mathfrak{m}}$ be a finite dimensional D weight module with support contained in orbit ω. Then $dim(V_{\mathfrak{m}}) = dim(V_{\mathfrak{n}})$ for any $\mathfrak{m}, \mathfrak{n} \in \omega$.*

Proof. Since the characteristic of $\Bbbk$ is p and so is the order of q, we see that ω is a circular orbit of length p. Further, ω has at most one A_q-break and has at most one A_1-break.

Suppose that ω does not have an A_q-break or an A_1-break. Let $\mathfrak{m} \in \omega$ be in the support of V, and consider $X : V_{\mathfrak{m}} \to V_{\alpha(\mathfrak{m})}$. Suppose $v \in V_{\mathfrak{m}}$ is in the kernel of X. That is, $YX(v) = \tau(v) = 0$ and $Y_1X(v) = (q\sigma - 1)(v) = 0$. That is, $\mathfrak{m} = (\tau, \sigma - \frac{1}{q})$ which implies that $\mathfrak{m}$ is an A_q as well as an A_1 break. That is, X is injective for every $\mathfrak{m}$. Since ω is a circular orbit, $X : V_{\mathfrak{m}} \to V_{\alpha(\mathfrak{m})}$ is an isomorphism of vectorspaces for every $\mathfrak{m}$.

Now suppose that ω has an A_q-break or an A_1-break. Without loss of generality, suppose that ω has an A_q-break, and since it is unique, let us name it $\mathfrak{m}_{p-1}$. Let $\omega = \{\mathfrak{m}_0, \mathfrak{m}_1, \ldots, \mathfrak{m}_{p-1}\}$ where $\mathfrak{m}_i = \alpha^i(\mathfrak{m}_0)$. As $\mathfrak{m}_i$ is not a break for any $i < p - 1$, then by the argument in the preceding paragraph, $X : V_{\mathfrak{m}_i} \to V_{\mathfrak{m}_{i+1}}$ is injective for every $i < p - 1$.

Suppose for some $i < p - 1$, $X : V_{\mathfrak{m}_i} \to V_{\mathfrak{m}_{i+1}}$ is not surjective. Then there is a $v \in V_{\mathfrak{m}_{i+1}}$ not in $X(V_{\mathfrak{m}_i})$. Now, $Y_1(v) \in V_{\mathfrak{m}_i}$. Note, $XY_1(v) = (\sigma - 1)(v)$. If $(\sigma - 1)(v) \neq 0$, then $v \in X(V_{\mathfrak{m}_i})$ contradicting the choice of v. Hence, $(\sigma - 1)(v) = 0$. In other words, $v \in \mathfrak{m}_0$. In other words, $X : V_{\mathfrak{m}_i} \to V_{\mathfrak{m}_{i+1}}$ is surjective for every $i, 0 \leq i < p - 1$. We have thus proved the theorem. ∎

Remark 14: *Note that by Proposition 1, every weight space of an irreducible D-module is of dimension 1 over an algebraically closed field, irrespective of its characteristic. Theorem 6 seems to lead to a generalization of Proposition 1 for D-modules not necessarily irreducible. Yet, Theorem 6 is not true for weight D-modules when the characteristic $\Bbbk$ is not equal to the order of q. For instance, suppose characteristic of $\Bbbk$ is 3 and $q^2 = 1$. Then, a cyclic orbit is of length 6. Let $\mathfrak{m}_1 = (\tau - 1, \sigma - 1)$ and $\omega = \{\mathfrak{m}_i = \alpha^i(\mathfrak{m}_1) \mid 0 \leq i \leq 5\}$. Let $V_{\mathfrak{m}_i}$ be one-dimensional with basis $\{v_i\}$ for $1 \leq i \leq 3$ and zero-dimensional otherwise. Assign it a D-module structure by*

$$\begin{aligned}
\tau(v_1) &= v_1, & \sigma(v_1) &= v_1, & X(v_1) &= v_2,\\
\tau(v_2) &= 2v_2, & \sigma(v_2) &= qv_2, & X(v_2) &= v_3,\\
\tau(v_3) &= 0, & \sigma(v_3) &= v_3, & X(v_3) &= 0,
\end{aligned}$$

$$\begin{aligned} Y(v_1) &= 0, & Y_1(v_1) &= 0, \\ Y(v_2) &= v_1, & Y_1(v_2) &= (q-1)v_1, \\ Y(v_3) &= 2v_2, & Y_1(v_3) &= 0. \end{aligned}$$

4.5. ISOMORPHISMS AMONG VARIOUS FAMILIES OF D-IRREDUCIBLE WEIGHT MODULES. The following isomorphisms are noted:

(1) When $b \in \Bbbk^* \setminus \{q^i\}_{i\in\mathbb{Z}}$ and $a \in \Bbbk \setminus \mathbb{Z}$, then $V_q(\omega, b, a) \cong V_1(\omega, a, b)$. These modules are described in section 3.1, case 3.1.1, and section 3.2, case 3.2.1, respectively.

(2) When q is not a root of unity, $b = q^k$ for some $k \in Z$ and $a \in \Bbbk \setminus \mathbb{Z}$, then $V_1(\omega, a, b) \cong V_q(\omega, J, J'a)$ where the two are described in section 3.2, case 3.2.1, and section 3.1, case 3.1.2, subcase (1) respectively.

(3) When q is a root of unity, $b = q^k$ for some $k \in Z$ and $a \in \Bbbk \setminus \mathbb{Z}$, then $V_1(\omega, a, b) \cong V_q(\omega, J, J)$ where the two are described in section 3.2, case 3.2.1, and section 3.1, case 3.1.3, case (1) respectively.

(4) When $b \in \Bbbk^* \setminus \{q^i\}_{i\in\mathbb{Z}}$ and $a \in \mathbb{Z}$, then $V_q(\omega, b, a) \cong V_1(\omega, J, J', b)$. These modules are described in section 3.1, case 3.1.1, and section 3.2, case 3.2.2, subcase (1) respectively.

(5) When $a \notin \mathbb{Z}_p$ and $q \notin \{q^i\}_{i\in\mathbb{Z}}$, we have $V_q(\omega, f, b, a) \cong V_1(\omega, f, a, b)$.

(6) When $w = \epsilon$, the emtpy word, $V_q(\omega, j, \epsilon)$ is isomorphic to $V_1(\omega, j, \epsilon)$ as D-modules; refer to sections 3.1.5 and 3.2.5 for details.

Acknowledgements

V.F. is supported in part by the CNPq grant (301320/2013-6) and by the Fapesp grant (2010/50347-9). U.I. gratefully acknowledges hospitality from Insitute of Mathematics and Statistics, University of São Paulo. She is supported in part by the Fapesp grant (2013/13970-8).

References

1. V. Bekkert, V., Benkart, G., Futorny, V., *Weight modules for Weyl algebras*, in "Kac-Moody Lie algebras and related topics (Proc. Ramanujan internat. symp., Chennai, India, 2002)", Contemp. Math. 343, Amer. Math. Soc., Providence, RI, (2004) 17-42.
2. Drozd, Yu. A., Guzner, B. L., Ovsienko, S. A., *Weight modules over generalized Weyl algebras,* J. Algebra 184 (1996) 491-504.
3. Iyer, U. N., Jordan, D. A., *Noetherian algebras of quantum differential operators*, (Work in progress)

4. Iyer, U. N., McCune, T. C., *Quantum differential operators on* $\Bbbk[x]$, Internat. J. of Math., Vol. 13, No.4 (2002) 395-413.
5. Lunts, V., Rosenberg, A., *Differential operators on noncommutative rings*, Selecta Math.(N.S) **3**, 335–359 (1997).